



Original Research Article

Incidence of Oral malignancy in Basrah Southern Iraq between 2005-2017

Sundus Abdul Wadood Algazaeri^{1,*}, Ghaydaa Hashim Al Qudsi²,
Hanaa Khazaal Jaber¹, Wasson Mansour Al Elwi³

¹Dept. of Oral and Maxillofacial Surgery, Basrah Dental College/University of Basrah, Basrah, Iraq

²Dept. of Oral Dignosis, Basrah Dental College/University of Basrah, Basrah, Iraq

³Al-Sadder Teaching Hospital, Basrah, Iraq



ARTICLE INFO

Article history:

Received 08-05-2020

Accepted 11.05.2020

Available online 21-07-2020

Keywords:

Oral malignancy

Squamous cell carcinoma

Incidence

Basrah

2005-2017

ABSTRACT

Background: Cancer is a major health problem in Basrah. Oral cancer is an important component which deserves exploration.

Objective: To study the incidence of oral malignancy in Basrah.

Materials and Methods: Available data were obtained from the cancer registry in Basrah, a histopathological research facility of Al-Sadder Teaching Hospital, Al-Basrah General Hospital and College of Dentistry in Basrah.

Results: In Basrah from 2005 to 2017, there were 304 cases identified of oral malignancy in various sources, 152 cases for both male and female. Squamous cell carcinoma accounted for 82.2%. The Tongue is the most frequent site followed by buccal mucosa, lower jaw then the upper jaw.

Conclusion: No solid evidence exists regarding the incidence of oral malignancy in Basrah governorate and the present information represent only an attempt to measure the extent and characteristics of oral malignant diseases. Comprehensive cancer registration is highly required supported by accurate information on new cases and on cancer-related mortality. The current level of registered oral malignant diseases is likely to be underestimated.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

Oral malignancy is one of the worldwide diseases, which possesses the eighth position among various kinds known about diseases. Every year, the total number of oral malignant new cases surpasses 300.000 cases. In spite of the fact that the differing circulation of oral malignant occurrence around the world, the quantity of new cases shows an expansion in various countries. The mortality burden may be up to about 130.000 individuals globally each year.¹

In 2010, the World Health Organization (WHO) revealed an oral malignancy death pace of around 2 per 100,000 in the Middle East, which is lower than that announced in India and the United States.² Oral malignant diseases have

different aetiology epidemiology and survival rate.³

Little is known about oral malignancy in Basrah. The requirement for far-reaching information about malignant growth forms is obligatory to design and set up control programs for the regular oral disease which might be amiable to counteraction, early discovery and cure. This study is an attempt to describe in quantitative terms the extent of oral malignant diseases over the years 2005-2017.

2. Materials and Methods

An aggregate of 304 patients equally and by chance were distributed between females and males The cases were compiled from records available at the histopathological research facility of Al-Sadder Teaching Hospital, Al-Basrah General Hospital and College of Dentistry in Basrah. Additionally Cancer Registration Section at the

* Corresponding author.

E-mail address: sendes1966@gmail.com (S. A. W. Algazaeri).

Department of Pathology in the College of Medicine in Basrah. Was used as a source of additional cases of oral cancer. This investigation incorporated the period between 2005-2017. Information on each case included gender, age, histopathological diagnosis site of cancer and year of diagnosis. Cases of oral malignancy whose normal residence is governorates other than Basrah were excluded. Sites of oral malignant growth included: upper and lower lip, tongue, gingiva, the floor of mouth, palate, buccal mucosa, upper and lower jaw Data were entered into an SPSS version 20. Population figures for Basrah population were obtained from various sources verified by the estimates published by the Ministry of planning and developmental cooperation.

Table 1: Time trend of incident oral malignancy in Basrah over the years 2005-2017

Year	Total population	No. of new cases	IR/100000
2005	2 173 197	15	0.69
2006	2 201 313	20	0.91
2007	2 267 352	21	0.93
2008	2 335 373	29	1.24
2009	2 405 434	35	1.46
2010	2 467 531	35	1.42
2011	2 531 997	42	1.66
2012	2 602 983	10	0.38
2013	2 675 774	19	0.71
2014	2 750 696	20	0.73
2015	2 827 715	21	0.74
2016	2 906 891	23	0.79
2017	3 071 956	14	0.45
Mid period population, number of cases and average IR/100000	2 531 997	23.4	0.92

3. Results

Among 304 cases analyzed as oral malignant diseases for the period 2005-2017, half of them were males and half were females. Oral malignancy is seen more among persons aged 50-59 years (23.7%), and this is true for both sexes (11.2%), for males and (12.5%) for females (Figure 1). Most cases were Squamous cell carcinoma (82.2%), females had slightly higher proportion of Squamous cell carcinoma 127(41.8%) than Males 123 (40.5%), then Non-Hodgkin s lymphoma about 12(3.9%), male 3.0% and female 1.0%, while different kinds of malignancies had little cases as found in (Figures 2, 3 and 4). There is a clear variation in the numbers and incidence rates of cases over the years. As shown in (Figure 5) and (Table 1).

4. Discussion

In Iraq and especially in Basrah in the southern part of Iraq, malignancy has certainly expanded in total incident cases, incidence rate and burden on the health care system.⁴⁻⁶

Most of the oral malignancy in this study was squamous cell carcinoma, like other going before researches, squamous cell carcinoma was the most widely recognized malignancy A great variation is reported in the mix of various types of malignancies among various studies^{2,3,7-13} in regard to the mix itself and/or in relation to sex.

As per the site, the research demonstrated the tongue as the most frequent site for an oral malignant growth in Basrah followed by buccal mucosa, a result which is consistent with to the pattern in European and the US populaces,^{2,14} but disagree with a study in Sulaimani⁸ that showed the lip as the first then the tongue. While the investigations among Asian populaces uncovered that the most widely recognized site is Buccal mucosa because of betel quid/tobacco biting habits.¹⁴ Remarkably, the gingiva and alveolus were the most ordinarily influenced locales among patients from the Southwestern district of Saudi Arabia (to be specific, Jizan and Najran).¹⁵ Besides, in Sudan, a high introduction of cancer that surpassed one anatomical region to attack the nearby one was observed.¹⁶

As per the age distribution, just 7.2% of malignancy happened in persons aged less than 30 years. The event of oral squamous cell carcinoma is uncommon among youthful people. The peak beginning of oral malignancy starts from the sixth decade onwards. This outcome might be perceived because of versatile resistant reactions in old people,^{10-12,15,16} and extensive stretch of exposure to various cancer-causing agents, for example, synthetic compounds, radiation and infections which are significant hazard factors causes oral malignancy.^{3,8-10,17,18} Alcohol consumption is viewed as a hazard factor for oral malignancy with cigarette smoking in Iraq, Libya and Egypt,² Utilization of Shamma and Qat were identified with the expanded danger of oral malignancy in Saudi Arabia and Yemen,^{12,13} but these are not prevailing in Basrah.

In Basrah an expanded danger of malignancy as an impression of expanded presentation to different hazard factors, for example, way of life changes, the aftermaths of different wars contamination and financial assents and different exposures are likely to be all operating to enhance the risk of malignant diseases.¹⁹

Most investigations don't record oral malignant growth like the investigation of disease in Basrah in a seven-year exertion throughout the years 2005-2011 made by the Basrah Cancer Research Group (BCRG) brought about enrolling an aggregate of 17080 new cases. Likewise Iraqi Cancer Board in Ministry Of Health every year distributed detailed measurable information of malignancy in Iraq, this incorporates the most widely recognized ten malignant diseases and appropriation of all tumours in




Fig. 1: Age group distribution in relation to sex




Fig. 2: Distribution of malignant types in relation to sex




Fig. 3: Site distribution in relation to sex




Fig. 4: Distribution of the malignancy according to the site




Fig. 5: Distribution of cases of oral malignancy according to years of diagnosis

various governorates and ages and genders with little data of oral malignancies.^{19,20} Concentrate International Agency for Research on Cancer/WHO in Iraq record just 0.98% of oral malignancy among 25.320 diseases recorded in 2018.²¹ In this investigation oral malignant growth recorded was 304 case, among them 209 cases recorded by Iraqi Cancer Registry in Basrah, that's means around 95 cases recorded from the different other sources. The implication is that official cancer registries are still lagging behind complete coverage of case detection and registration. In this context, our results support the conclusions stated by Al-Hilfe and Habib in 2016 when they reported an under-registration of around one-fifth of the incident cancer cases in Basrah.²² Certainly, under-registration as well as expanding the scope of Iraqi Cancer Board are significant future challenges facing this institution.²³

5. Conclusion

No solid evidence exists regarding the incidence of oral malignancy in Basrah governorate and the present information not very identical with the Iraqi cancer registry, this registry must include more information about the risk factor, address, habits, economic, treatment pattern and mortality to all types of malignancies included oral

malignancy on each province. High-quality, accurate data based oral malignant registries, with accurate mortality statistics, serve to limit the unreported and undiagnosed oral malignancy cases and help better guides health policy pioneers in building up proper diagnostic, treatment and preventive services all through Iraq. Ministry of Health in Iraq must search for a system to confirm the medical conditions that its patients wish to receive treatment outside the country and the reasons for which the patient desires treatment abroad and try to find solutions to a healthy reality appropriate for the Iraqi people.

6. Source of Funding

None.

7. Conflict of Interest

None.

References

1. Khattab NMA, Elheeny AAH, Tony GA. Oral-cancer knowledge, practice, and attitude assessment of dentists in Upper Egypt: A cross-sectional study. *Clin Exp Dent Res*. 2019;5(2):121–7.
2. Al-Jaber A, Al-Nasser L, El-Metwally A. Epidemiology of oral cancer in Arab countries. *Saudi Med J*. 2016;37(3):249–55.

3. Zini A, Czerninski R, Sgan-Cohen HD. Oral cancer over four decades: epidemiology, trends, histology, and survival by anatomical sites. *J Oral Pathol Med*. 2009;39(4):299–305.
4. Habib OS, Hameed LA, Ajeel NA, Al-Hawaz MH, Al-Faddagh ZA, Nasr GN, et al. Epidemiology of Breast Cancer among Females in Basrah. *Asian Pac J Cancer Prev*. 2016;17(sup3):191–5.
5. A OSH, A KAAI. A changing pattern of cancer-related mortality in Basrah. *Basrah J Surg*. 2018;24(2):24–9.
6. Habib OAE. Pattern of Cancer: Comparison of risk in Basrah with national pattern. *Med J Babylon*. 2018;15:195–6.
7. BenNasir E, Mistiri ME, McGowan R, Katz RV. Oral cancer in Libya and development of regional oral cancer registries: A review. *Saudi Dent J*. 2015;27(4):171–9.
8. Talabani NG, Ahmed KM, Faraj FH, and. Oral Cancer in Sulaimani: A Clinicopathological Study. *J Zankoy Sulaimani - Part A*. 2008;13(1):1–8.
9. Museedi OS, Younis WH. Oral cancer trends in Iraq from. *Saudi J Dent Res*. 2000;5:41–7.
10. Mm AM, Alsaimry IE, Al AA. Occurrence of oral and oropharyngeal squamous cell carcinoma among patients in Basrah city. 2017;16:255–61.
11. Gundamaraju KK, Kantheti LPC, Naga SNDV, Navakoti P, Poosarla C, Bujunuru SR. Prevalence of oral potentially malignant and malignant lesions at a tertiary level hospital in Hyderabad, India. *J Dr NTR Univ Heal Sci*. 2014;3(5):13.
12. Brown A, Ravichandran K, Warnakulasuriya S. The unequal burden related to the risk of oral cancer in the different regions of the Kingdom of Saudi Arabia. *Community Dent Health*. 2006;23:101–6.
13. Sawair FA, Al-Mutwakel A, Al-Eryani K, Al-Surhy A, Maruyama S, Cheng J, et al. High relative frequency of oral squamous cell carcinoma in Yemen: Qat and tobacco chewing as its aetiological background. *Int J Environ Health Res*. 2007;17(3):185–95.
14. Warnakulasuriya S. Global epidemiology of oral and Oropharyngeal cancer. *Oral Oncol*. 2009;45:309–16.
15. Basha I S, Mohamed RN, Ya-T, Asas. The Prevalence of Oral Cancer in Saudi Arabia - A Systematic Review. *Ann Med Heal Sci Res*. 2019;9:553–7.
16. Osman TA, Satti AA, Bøe OE. Pattern of malignant tumors registered at a referral oral and maxillofacial hospital in Sudan during. *J Cancer Res Ther*. 2006;6:473–7.
17. Zaib N. Oral biopsies : a study of 114 cases ORAL BIOPSIES : STUDY OF 114 CASES. *Pakistan Oral Dent. J*. 2012;32:416–20.
18. Medicine O, Perriman BDS, DHD, S FD. Department of Oral Surgery and Oral Medicine, College of Dentistry BU. ORAL CANCER IN IRAQ: ANALYSIS OF 202 CASES. *Br J Oral Surg*. 1964;179:146–51.
19. Hussain RA, Habib OS. Incidence of Cancer in Basrah : Results of a Household Survey Incidence of Cancer in Basrah : Results of a Household Survey. *APJCP*. 2015;16:163.
20. Health MOF. Republic of iraq ministry of health iraqi cancer board Iraqi cancer registry. Baghdad: Republic of iraq ministry of health; 2015.
21. Country-specific I, Method N, Country-specific M. Iraq source:Globocan 2018. *Int Agency Res Cancer/WHO*. 2018;141:2018–9.
22. Abdul-Ameer R, Habib OS. Cancer Registration in Basrah-Southern Iraq: Validation by household survey. *Asian Pacific J Cancer Prev*. 2016;17(3):197–200.
23. Partow Z, Kulaksiz BGMS, Goonesekere ZBR. Report No . 112333-IQ Systematic Country Diagnostic; 2017.

Author biography

Sundus Abdul Wadood Algazaeri Lecturer Oral Surgeon

Ghaydaa Hashim Al Qudsi Lecturer Oral Pathology

Hanaa Khazaal Jaber Consultant Histopathologist

Wasson Mansour Al Elwi Consultant Histopathologist

Cite this article: Algazaeri SAW, Al Qudsi GH, Jaber HK, Al Elwi WM. Incidence of Oral malignancy in Basrah Southern Iraq between 2005-2017. *Int J Oral Health Dent* 2020;6(2):110-115.