

## Knowledge, Awareness and Practices regarding Sharp Injuries amongst the Dental students

Rana Pratap Maurya<sup>1</sup>, Malti Kumari Maurya<sup>2,\*</sup>, Rashmi Kushwaha<sup>3</sup>, Sneha Lata Verma<sup>4</sup>, Reema Kumari<sup>5</sup>

<sup>1,4</sup>Reader, Dept. of Orthodontics & Dentofacial Orthopaedics, Babu Banarasi Das College of Dental Sciences, Lucknow, Uttar Pradesh, <sup>2,3</sup>Associate Professor, Dept. of Pathology, <sup>5</sup>Professor, Dept. of Community Medicine, King George's Medical University, Lucknow, Uttar Pradesh

**\*Corresponding Author:**

Email: mauryamalti@yahoo.co.in

### Abstract

**Introduction:** Dental students are at risk to various blood borne diseases transmitted by needle stick/sharp injuries (NSIs) during the course of their clinical training which have been. To reduce the risk from NSIs there should be an essential need to improve and update the knowledge of NSIs and its management by continued education programmes as seminars, lectures and workshops.

**Aim:** To assess the knowledge, awareness and practices regarding sharp injuries amongst the dental students.

**Materials and Method:** This cross-sectional survey based study was conducted among the 150 voluntarily participated dental students who were receiving their undergraduate clinical training in various dental colleges. Data was recorded on a structured questionnaire to elicit the level of knowledge, awareness and practice towards sharp injuries. All data was tabulated in terms of number and percentage.

**Results:** 64% dental students experience NSI in last six month. The main cause of NSIs was orthodontic wires (18.75%) followed by explorer (15.63%), endodontic file and scalpel blade (14.58%), scaler (11.46%), hollow needle and bur (9.38%). Highest number of NSIs was found during orthodontic wire bending and treatment (18.75%) followed by the endodontic treatment (15.63%), diagnosis (12.50%), scaling (10.41%), needle recapping and restoration (8.33%). Adequate number of students had good knowledge and awareness regarding Needle stick/sharp injury. In practice, maximum number of dental students washed hands, used glove and recapped needle after uses but only few students dispose sharps in puncture proof container and used needle destroyer before discarding the syringe during clinical procedure.

**Conclusion:** Dental students required training and teaching regarding management of sharp injuries and should be encouraged to report it to concerned authority.

**Keywords:** Needle stick/Sharp injury, Post exposure prophylaxis, Dental Students

### Introduction

Health care workers (HCWs) are least concerned for their own health however, they are at increased risk for acquiring more than 20 different pathogens due to occupational exposure to blood and body fluids.<sup>(1)</sup> Needle stick injury/ Sharp injury (NSI) is defined "as par lital introduction into body of health care worker, during the performance of their duties, of blood or other potentially hazardous material by hollow bore needle or sharp instruments, including, but not limited to needles, lancets, scalpels, and contaminated broken glass; constitute a major hazard for the transmission of various blood borne diseases such as Hepatitis-B, Hepatitis-C and HIV.<sup>(2)</sup> The risk of infection depends on the prevalence of disease in patient population, type of viral infection, viral load and the precautions taken by the HCWs while dealing these patients.

According to WHO, the annual estimated proportions of HCWs exposed to blood borne pathogens globally were 2.6% for HCV, 5.9% for HBV and 0.5 % for HIV.<sup>(3,4)</sup> More than 90% of these infections occur in developing countries but most of these NSIs remain unreported.<sup>(5)</sup> NSI results in psychological impacts as tension and fear and distraction from their work. Most of the needle stick injuries can be prevented by using safety

devices and by applying "Universal precautions" as a safety measure.

Dental students who works in various dental departments like, oral surgery, endodontics, orthodontics, prosthodontics, pedodontics and periodontics are generally at a higher risk of occupational hazard due to lack of experience and skill in performing dental procedures during clinical training periods.<sup>(6-7)</sup> In dental practice, various sharp instruments and syringes are used that pose them at increased risk of sustaining NSIs.<sup>(8-9)</sup>

Despite the increased rate of NSIs among practicing dentists, there is no research on needle stick/sharp injuries among the undergraduate dental student in Lucknow. Hence, the aim of the present study was to assess the knowledge, awareness and practices regarding sharp injuries amongst the undergraduate and intern dental students in Lucknow, India.

### Materials and Method

This cross-sectional study was conducted among the 150 dental students of B.D.S third year, final year and interns (50 each), from the various dental colleges of Lucknow, India. All subjects had voluntarily participated in the study and were fully informed about the design and purpose of the study.

A questionnaire was prepared and distributed among the dental students to elicit the level of knowledge, awareness and practice towards sharp injuries. All data was tabulated in terms of number and percentage.

## Results

Results about knowledge, awareness and practices regarding NSIs among overall dental students and for different batches of dental students are shown in Table 1 and 2. The main cause of NSIs among all dental students was orthodontic wires (18.75%) followed by explorer (15.63%), endodontic file and scalpel blade (14.58%), scaler (11.46%), hollow needle, bur (9.38%) and extraction instrument (6.25%). Maximum sharp injuries in BDS 3<sup>rd</sup> students was due to wires during

Orthodontic treatment, for BDS final year student by scalpel blade while for interns by endodontic files during Endodontic treatment (Table 3 and Fig. 1). Procedures during which highest number of NSIs occurs was orthodontic wire bending and treatment (18.75%) followed by the endodontic treatment (15.63%), diagnosis (12.50%), scaling (10.41%), needle recapping and restoration (8.33%), washing of sharp instruments and disposal of sharp instruments (7.29%), extraction/surgical procedure (5.20%), collision with sharp instrument (4.17%) and least during local anesthesia administration (2.08%). In BDS 3<sup>rd</sup> year and final year student highest number of NSIs were observed during orthodontic wire bending/treatment while in interns it was found during endodontic treatment (Table 4 and Fig. 2).

**Table 1: Knowledge, Awareness and Practice regarding sharp/needle stick injury (NSIs) among overall dental students**

| S.N.   | Questions about Knowledge and Awareness regarding NSIs                              | Overall Dental Students (N=150) |                 |       |
|--|---|---------------------------------|-----------------|-------|
|  |   | Number (N)                      | Percentages (%) |       |
| 1.   | Definition of Needle stick/Sharp injury (NSIs)                                      | 105                             | 70.00           |       |
| 2.   | Risk of NSIs  | 132                             | 88.00           |       |
| 3.   | More than 20 different types of pathogen may be transmitted by NSIs                 | 125                             | 83.33           |       |
| 4.   | Hepatitis-B can be prevented by vaccine   | 142                             | 94.67           |       |
| 5.   | Currently no vaccine is present to prevent Hepatitis-C infection                    | 105                             | 70.00           |       |
| 6.   | Knowledge about universal precaution  | 123                             | 82.00           |       |
| 7.   | NSI is an occupational hazard in dental community                                   | 123                             | 82.00           |       |
| 8.   | Hypodermic needle increases the risk for NSIs                                       | 116                             | 77.33           |       |
| 9.   | Most NSIs occurred during recapping of used needle                                  | 122                             | 81.33           |       |
| 10.  | Most of NSIs have been neglected and unreported                                     | 106                             | 70.67           |       |
| 11.  | Affected area should be rinsed and washed thoroughly with soap and water after NSIs | 140                             | 93.33           |       |
| 12.  | Post exposure prophylaxis (PEP)   | 110                             | 73.33           |       |
| 13.  | Timing of Post-exposure Prophylaxis   | 102                             | 68.00           |       |
| 14.  | Stress and anxiety after NSIs   | 106                             | 70.67           |       |
| 15.  | Hazardous nature of biomedical waste  | 110                             | 73.33           |       |
| 16.  | Proper hand washes before and after procedure                                       | 128                             | 85.33           |       |
| 17.  | Most NSIs occurred during disposal of used needles                                  | 121                             | 80.67           |       |
| 18.  | Segregation of biomedical waste at the point of its generation                      | 122                             | 81.33           |       |
| 19.  | Universal precaution  | 124                             | 82.67           |       |
| 20.  | Personal preventive measure   | 111                             | 74.00           |       |
| 21.  | Correct method of sharp disposal  | 95                              | 43.33           |       |
| 22.  | Recapping of needle after use   | 120                             | 80.00           |       |
| 23.  | Reporting of needle stick injury  | 101                             | 67.33           |       |
| <b>Questions about Practice regarding NSIs</b> |   |                                 |                 |       |
| 24.  | Dealing with any sharp objects during clinical training                             | 148                             | 98.67           |       |
| 25.  | Experience about any NSI in last six month  | 96                              | 64.00           |       |
| 26.  | Number of NSIs in last 6 months   | 1-2 times                       | 63              | 65.62 |
|  |   | 3-4 times                       | 24              | 25.00 |
|  |   | More than 4 times               | 9               | 9.38  |
| 27.  | Injury washed with soap and water immediately                                       | 88                              | 91.67           |       |
| 28.  | NSIs reported to the assigned authority   | 53                              | 55.20           |       |
| 29.  | Received any teaching/training in BMW   | 53                              | 35.33           |       |

|     |  |     |       |
|-----|--|-----|-------|
| 30. | Disposal of all kinds of waste including sharps into general garbage       | 132 | 88.00 |
| 31. | Recapping of needle after its use  | 122 | 81.33 |
| 32. | Dispose needle and sharp objects after its use in puncture proof container | 10  | 6.67  |
| 33. | Wear gloves and other protective measures                                  | 112 | 74.67 |
| 34. | Washed hands before and after procedure                                    | 125 | 83.33 |
| 35. | Used needle destroyer before discarding needles                            | 8   | 5.33  |
| 36. | Vaccinated for Hepatitis-B   | 145 | 96.67 |




Fig. 1: Cause and incidence of sharp/ needle stick injury (NSIs) among Dental Students

Table 2: Knowledge, Awareness and Practice regarding sharp/ needle stick injury (NSIs) among different batches of dental students

| S.N. | Questions about Knowledge and Awareness regarding NSIs                              | Third Year (N=50) |       | Final Year (N=50) |       | Intern (N=50) |       |
|------|---|-------------------|-------|-------------------|-------|---------------|-------|
|      |   | N                 | %     | N                 | %     | N             | %     |
| 1.   | Definition of Needle stick/Sharp injury (NSIs)                                      | 31                | 62.00 | 35                | 70.00 | 39            | 78.00 |
| 2.   | Risk of NSIs  | 42                | 84.00 | 44                | 88.00 | 46            | 92.00 |
| 3.   | More than 20 different types of pathogen may be transmitted by NSIs                 | 39                | 78.00 | 41                | 82.00 | 45            | 90.00 |
| 4.   | Hepatitis-B can be prevented by vaccine   | 40                | 80.00 | 45                | 90.00 | 47            | 94.00 |
| 5.   | Currently no vaccine is present to prevent Hepatitis-C infection                    | 30                | 60.00 | 35                | 70.00 | 40            | 40.00 |
| 6.   | Knowledge about universal precaution  | 35                | 70.00 | 42                | 84.00 | 46            | 92.00 |
| 7.   | NSI is an occupational hazard in dental community                                   | 38                | 76.00 | 41                | 82.00 | 44            | 88.00 |
| 8.   | Hypodermic needle increases the risk for NSIs                                       | 37                | 74.00 | 39                | 78.00 | 40            | 80.00 |
| 9.   | Most NSIs occurred during recapping of used needle                                  | 40                | 80.00 | 41                | 82.00 | 41            | 82.00 |
| 10.  | Most of NSIs have been neglected and unreported                                     | 35                | 70.00 | 32                | 64.00 | 39            | 78.00 |
| 11.  | Affected area should be rinsed and washed thoroughly with soap and water after NSIs | 45                | 90.00 | 47                | 94.00 | 48            | 96.00 |
| 12.  | Post exposure prophylaxis (PEP)   | 34                | 68.00 | 38                | 76.00 | 38            | 76.00 |
| 13.  | Timing of Post-exposure Prophylaxis   | 30                | 60.00 | 35                | 70.00 | 37            | 74.00 |
| 14.  | Stress and anxiety after NSIs   | 35                | 70.00 | 34                | 68.00 | 37            | 74.00 |
| 15.  | Hazardous nature of biomedical waste  | 32                | 64.00 | 36                | 72.00 | 42            | 84.00 |
| 16.  | Proper hand washes before and after procedure                                       | 40                | 80.00 | 43                | 86.00 | 45            | 90.00 |
| 17.  | Most NSIs occurred during disposal of used needles                                  | 39                | 78.00 | 41                | 82.00 | 41            | 82.00 |
| 18.  | Segregation of biomedical waste at the point of its generation                      | 36                | 72.00 | 42                | 84.00 | 44            | 88.00 |
| 19.  | Universal precaution  | 37                | 74.00 | 42                | 84.00 | 45            | 90.00 |

|  |  |                   |        |       |        |       |        |       |
|--|--|-------------------|--------|-------|--------|-------|--------|-------|
| 20.  | Personal preventive measure  | 34                | 68.00  | 39    | 78.00  | 38    | 76.00  |       |
| 21.  | Correct method of sharp disposal   | 30                | 60.00  | 32    | 64.00  | 33    | 66.00  |       |
| 22.  | Recapping of needle after use  | 37                | 74.00  | 40    | 80.00  | 43    | 86.00  |       |
| 23.  | Reporting of needle stick injury   | 30                | 60.00  | 35    | 70.00  | 36    | 72.00  |       |
| <b>Questions about Practice regarding NSIs</b> |  |                   |        |       |        |       |        |       |
| 24.  | Dealing with any sharp objects during clinical training                    | 48                | 96.00  | 50    | 100.00 | 50    | 100.00 |       |
| 25.  | Experience about any NSI in last six month                                 | 28                | 56.00  | 36    | 72.00  | 32    | 66.00  |       |
| 26.  | Number of NSIs in last 6 months  | 1-2 times         | 18     | 64.29 | 23     | 46.00 | 22     | 68.75 |
|  |  | 3-4 times         | 7      | 25.00 | 9      | 18.00 | 8      | 25.00 |
|  |  | More than 4 times | 3      | 10.71 | 4      | 8.00  | 2      | 6.25  |
| 27.  | Injury washed with soap and water immediately                              | 25                | 89.29  | 34    | 94.44  | 29    | 90.63  |       |
| 28.  | NSIs reported to the assigned authority                                    | 15                | 53.58  | 21    | 58.33  | 17    | 53.12  |       |
| 29.  | Received any teaching/training in BMW                                      | 13                | 26.00  | 19    | 38.00  | 21    | 42.00  |       |
| 30.  | Disposal of all kinds of waste including sharps into general garbage       | 43                | 86.00  | 45    | 90.00  | 44    | 88.00  |       |
| 31.  | Recapping of needle after its use  | 38                | 76.00  | 43    | 46.00  | 41    | 82.00  |       |
| 32.  | Dispose needle and sharp objects after its use in puncture proof container | 4                 | 8.00   | 3     | 6.00   | 3     | 6.00   |       |
| 33.  | Wear gloves and other protective measures                                  | 34                | 68.00  | 37    | 74.00  | 41    | 82.00  |       |
| 34.  | Washed hands before and after procedure                                    | 39                | 78.00  | 41    | 82.00  | 45    | 90.00  |       |
| 35.  | Used needle destroyer before discarding needles                            | 3                 | 6.00   | 2     | 4.00   | 3     | 6.00   |       |
| 36.  | Vaccinated for Hepatitis-B   | 50                | 100.00 | 48    | 96.00  | 47    | 94.00  |       |




Fig. 2: Sharp/ Needle sticks injury (NSIs) among Dental Students during various Clinical procedures

Table 3: Cause and incidence of sharp/ needle stick injury (NSIs) among dental students

| S. N. | Cause of NSIs         | Third Year (N=28) |       | Final Year (N=36) |       | Intern (N=32) |       | Total (N=96) |       |
|-------|-----------------------|-------------------|-------|-------------------|-------|---------------|-------|--------------|-------|
|       |                       | N                 | %     | N                 | %     | N             | %     | N            | %     |
| 1.    | Hollow Needle         | 1                 | 3.57  | 3                 | 8.33  | 5             | 15.62 | 9            | 9.38  |
| 2.    | Explorer              | 6                 | 21.42 | 5                 | 13.89 | 4             | 12.50 | 15           | 15.63 |
| 3.    | Endodontic File       | 1                 | 3.57  | 5                 | 13.89 | 8             | 25.00 | 14           | 14.58 |
| 4.    | Scalpel blade         | 5                 | 17.86 | 7                 | 19.44 | 2             | 6.25  | 14           | 14.58 |
| 5.    | Scaler                | 5                 | 17.86 | 3                 | 8.33  | 3             | 9.38  | 11           | 11.46 |
| 6.    | Extraction Instrument | 0                 | 0.00  | 3                 | 8.33  | 3             | 9.38  | 6            | 6.25  |
| 7.    | Orthodontic Wire      | 8                 | 28.57 | 6                 | 16.67 | 4             | 12.50 | 18           | 18.75 |
| 8.    | Bur                   | 2                 | 7.14  | 4                 | 11.11 | 3             | 9.38  | 9            | 9.38  |

Table 4: Sharp/ needle sticks injury (NSIs) among dental students during various clinical procedures

| S. N. | Clinical Procedure                  | Third Year (N=28) |       | Final Year (N=36) |       | Intern (N=32) |       | Total (N=96) |       |
|-------|-------------------------------------|-------------------|-------|-------------------|-------|---------------|-------|--------------|-------|
|       |                                     | N                 | %     | N                 | %     | N             | %     | N            | %     |
| 1.    | Local anaesthesia administration    | 0                 | 0.00  | 1                 | 2.78  | 1             | 3.13  | 2            | 2.08  |
| 2.    | Diagnosis                           | 5                 | 17.86 | 4                 | 11.11 | 3             | 9.38  | 12           | 12.50 |
| 3.    | Endodontic treatment                | 1                 | 3.57  | 5                 | 13.89 | 9             | 28.13 | 15           | 15.63 |
| 4.    | Extraction/ Surgical procedure      | 0                 | 0.00  | 3                 | 8.33  | 2             | 6.25  | 5            | 5.20  |
| 5.    | Scaling                             | 4                 | 14.28 | 3                 | 8.33  | 3             | 9.38  | 10           | 10.41 |
| 6.    | Needle recapping                    | 1                 | 3.57  | 4                 | 11.11 | 3             | 9.38  | 8            | 8.33  |
| 7.    | Collision with sharp instrument     | 1                 | 3.57  | 2                 | 5.56  | 1             | 3.13  | 4            | 4.17  |
| 8.    | Orthodontic wire bending/ Treatment | 8                 | 28.57 | 6                 | 16.67 | 4             | 12.50 | 18           | 18.75 |
| 9.    | Restoration                         | 2                 | 7.14  | 3                 | 8.33  | 3             | 9.38  | 8            | 8.33  |
| 10.   | Washing of sharp instruments        | 3                 | 10.71 | 3                 | 8.33  | 1             | 3.13  | 7            | 7.29  |
| 11.   | Disposal of sharp instruments       | 3                 | 10.71 | 2                 | 5.56  | 2             | 6.25  | 7            | 7.29  |

## Discussion

In the present study, most of the students were familiar with the risk of NSI during clinical procedures and they knew the definition of NSIs. They also had knowledge about pathogens transmitted by sharp injuries, vaccine available for Hepatitis Band non-availability of hepatitis-C vaccine. Similar findings were found by Saini R<sup>(10)</sup> Gichki et al<sup>(11)</sup> and Lakshmi et al<sup>(12)</sup> in their study. Whereas, Garima et al<sup>(13)</sup> in their study reported that only 22% students knew the definition of NSIs and 58.6% knew the immediate measures to be taken.

Regarding the causes of sharp injuries and the universal precautions most of the students were aware that the major sharp injuries occurred during recapping of used needle. 70.67% students were agreed that most of NSIs had been neglected and unreported while 93.33% was aware that after sharp injuries, affected area should be rinsed and washed thoroughly with soap and water. Most of the students knew the importance of post exposure prophylaxis (PEP) and correct timing for post-exposure prophylaxis. In contrast to our findings Cervini and Bell,<sup>(14)</sup> Alam M<sup>(15)</sup> and Siddiqui et al<sup>(16)</sup> found PEP practices for NSI were inadequate among medical students. Present study results showed that sharp injury may lead to significant stress and anxiety for the affected injured person whereas Saini et al<sup>(10)</sup> reported only 47% students lead to stress and anxiety after NSIs. Osman<sup>(17)</sup> found anxiety and stress the most frequent immediate post injury reaction reported by 59.4% students and 28.1% directed anger to themselves while 12.5% felt indifferent following the injury.

Maximum students were aware to ill effects of the biomedical waste and also believed that it should be segregated at the point of its generation so that risk of transmission of diseases can be reduced. 80% students believed that needle should be recapped after use and 67.33% students said that all needle stick injury should be reported but only 43% participants had knowledge that all the sharps should be disposed in puncture proof container to avoid injury.

Regarding practice and incidence about sharp injuries it was found that approximately all students deal with sharp objects during their daily clinical training. Amongst them 64% students experienced to NSIs during clinical training in last six month. Prevalence of NSI differed invariably by the different authors in their studies.<sup>(13-14,18-22)</sup>

In the present study maximum sharp injuries were caused by the orthodontic wires (18.75%) followed by explorer (15.63%), endodontic file and scalpel blade (14.58%), scaler (11.46%), hollow needle and bur (9.38%) and minimum by extraction instrument (6.25%) during respective procedures. In other studies Endodontics, surgery, prosthodontics, operative dentistry, pediatrics and periodontics departments were found as places for occurrence of injury.<sup>(10,15-16,26)</sup> Other procedures for injuries identified in this study such as sharps disposal, needle re-capping, washing instruments, scaling, wound suturing were also reported in similar studies.<sup>(6,8,18,21,23)</sup>

In the present study, students who get NSIs, 91.67% washed injured sites immediately with soap and water but and only 55.20% reported NSIs to the assigned authority. The action taken after NSIs in the study by Verma et al<sup>(19)</sup> included washing the site with soap and water, applying alcohol/betadine/antiseptics, expressing blood from injured site, applying pressure and tying the part. Sharma et al<sup>(21)</sup> reported that 60.9% HCWs washed the site of injury with water and soap and only 7.8% of the HCWs took post-exposure prophylaxis (PEP) against HIV/AIDS after NSIs. They also observed that patient overload and fatigue due to long hours of working was the commonest reason for causing the needle stick injury. Gichki et al<sup>(11)</sup> found that 99% respondents believed that the injury should be reported and 91% agreed that the post exposure prophylaxis should be initiated within one hour of injury. Lakshmi et al<sup>(12)</sup> in their study found that most of the students were aware NSI would have a psychological effect and diseases transmitted after an NSI can be prevented by prophylaxis and vaccinations.

In this study, only 35.33% had received teaching about biomedical waste management. Most of the students disposed all kinds of waste including sharps into general garbage and had practiced to recapped needle after its use. Very few students were used needle destroyer before discarding needles and disposed needle and sharp objects in puncture proof container. Most of the students used gloves and other protective measures during clinical procedure and washed hands before and after any clinical procedure. Lakshmi et al<sup>(12)</sup> reported that 64.1%, students were aware to reduce the occurrence of NSI by wearing gloves while 88.5% students were aware that NSI could transmit diseases transmitted by blood and 79.5% students knew that HIV and HBV could be transmitted by NSI. HBV exposures pose the highest risk for infection, but had an effective vaccine for it. Although Post exposure prophylaxis for health care workers can dramatically reduce the risk of infection but is not so effective for HCV and HIV. Approximately all students were vaccinated for hepatitis-B vaccine in present study. Other studies showed different range of vaccinations from ranging from 60% - 100%.<sup>(6,9,13,18,24-25)</sup> Unvaccinated individuals may have a 6%-30% chances of becoming infected with the virus following an injury.<sup>(6)</sup> However, the effectiveness of the vaccine is the most important factor that needs to be tested among healthcare students.<sup>(7)</sup>

### Conclusions

Sharp injuries can be reduced by following standard protocol regarding the training as well as adapting preventive measures. Reporting a sharp injury and the use of prophylactic measures are quite important. Therefore, every dental care centres and colleges should have infection control committee for providing infection control training and to look after the injured individuals.

### References

- Aiken LH, Sloane DM, Klocinski JL. Hospital nurses occupational exposure to blood: Prospective, retrospective, and institutional reports. *Am J Public Health* 1997; 87:103-07.
- Needle stick injuries. 2003 (cited January 23, 2008); Available from: <http://www.jr2.ox.ac.uk/bandolier/Extraforbando/needle.pdf>.
- Pruss-Ustun A, Rapiti E, Hutin Y. Sharps injuries: Global burden of diseases from sharps injuries to health-care workers. Geneva: World Health Organization; 2003.(WHO Environmental Burden of Diseases Series, No. 3).
- Tadesse M, Tadesse T. Epidemiology of needle stick injuries among health-care workers in Awassa City, Southern Ethiopia. *TROPICAL DOCTOR* 2010;40: 111-113 DOI: 10.1258/td.2009.090191.
- WHO. Reducing risks, promoting healthy life. Geneva; 2002. (cited 2013 Sep 7). Available from: [http://www.who.int/whr/2002/en/whr02\\_en.pdf](http://www.who.int/whr/2002/en/whr02_en.pdf).
- Jaber MA. A survey of needle sticks and other sharp injuries among dental undergraduate students. *Int J Infect Control* 2011; 7(3).
- Hussain JSA, Ram SM, Galinde J, Jingade RRR. Occupational exposure among dental, medical and nursing students in Mahatma Gandhi Mission's Campus, Navi Mumbai, India. *J Contemp Dent* 2012; 2(2).
- Gaballah K, Warbuton D, Sihmbly K, Renton T. Needle stick injuries among dental students: risk factors and recommendations for prevention. *Libyan J Med. Epub* 2012; 7 (1), 17507.
- McCarthy GM, Britton JE. A Survey of Final-Year Dental, Medical and Nursing Students: Occupational Injuries and Infection Control. *J Can Dent Assoc.* 2000 Nov;66(10):561
- Saini R. Knowledge and awareness of needle stick injury among students of Rural Dental College, Maharashtra, India. *Ann Nigerian Med* 2011; 5, (1):12-14.
- Gichki AS, Islam A, Murad W. Knowledge and awareness about needle stick injuries among dental students of Bolan Medical College, Quetta. *Pakistan Oral Dent J.* 2015;35(4):562-566.
- Lakshmi S, Bhat R. A Study on Awareness of Needle sticks injury in students undergoing paramedical course. *IOSR Journal of Dental and Medical Sciences IOSR-JDMS.* 2015, 14,(11)34-38.
- Garima M, Anmol T, Garwal R K, Gupta P, Gupta P. Knowledge, awareness and prevalence of needle stick injury among Students of medical college of Uttarakhand, India. *International Journal of Recent Scientific Research.* 2015,6(3):3055-3058.
- Cervini P, Bell C. Brief report: Needlestick injury and inadequate post-exposure practice in medical students. *J Gen Intern Med* 2005;20(5):419-21.
- Alam M. Knowledge, attitude and practices among health care workers on needlestick injuries. *Ann Saudi Med* 2002;22(5-6):396-9.
- Siddique K, Mirza S, Tauqir SF, Anwar I, Malik AZ. Knowledge attitude and practices regarding needlestick injuries amongst healthcare providers. *Pak J Surg* 2008;24(4):243-8.
- Osman T. Epidemiology of sharp instruments injuries at a dental school in Sudan. *Int J Infect Control* 2014, v10:i4.
- Askarian M, Malekmakan L, Memish ZA, Assadian O. Prevalence of needle stick injuries among dental, nursing and midwifery students in Shiraz, Iran. *GMS Krankenhhyg Interdiszip* 2012; 7(1): Doc05.
- Verma Y, Bandlish D, Kumar B. The prevalence of needle sticks injury among healthcare personnel in a tertiary care hospital in Kolkata. *Webmed Central Clinical Microbiology* 2015;6(9):WMC004983
- Gupta DK, Agrawal VK, Gupta SB, Ahmad Faheem: Needle Stick Injuries among Health Care Worker People's Journal of Scientific Research July 2015; Volume 8, Issue 2.
- Sharma R, Rasania S, Verma A, Singh S. Study of prevalence and response to needle stick injuries among health care workers in a tertiary care hospital in Delhi, India. *Indian J Community Medicine.* 2010;35: 74-77.
- Pavithran VK, Murali R, Krishna M, Shamala A, Yalamalli M, Kumar A V. Knowledge, attitude, and practice of needle stick and sharps injuries among dental professionals of Bangalore, India. *J Int Soc Prevent Community Dent* 2015;5: (5), 406-412.
- Hashemipour M, Sadeghi A. Needlestick injuries among medical and dental students at the University of Kerman. A questionnaire study. *J Dent Tehran Univ Med Scien Tehran Iran.* 2008; 5:71-76.
- Wicker S, Jung J, Allwinn R, Gottschalk R, Rabenau HF: Prevalence and prevention of needlestick injuries among

- health care workers in a German University hospital. *Int Arch Occup Environ Health.* 2008;81(3):347-354.
25. Radha R, Khan A. Epidemiology of Needle Stick Injuries Among The Health Care Workers of A Rural Tertiary Care Hospital-A Cross Sectional Study. *Natl J Community Med* 2012; 3(4):589-94.