

Content available at: <https://www.ipinnovative.com/open-access-journals>

International Journal of Oral Health Dentistry

Journal homepage: www.ijohd.org

Original Research Article

The distribution of malocclusion using the index of orthodontic treatment needs at a university dental hospital in and around Pretoria, South Africa

Tshepiso D Ntseke¹, Thomas K Madiba², Millicent Motsepe¹,
Mpule AL Moshaoa¹, Phumzile Hlongwa^{1*}

¹Dept. of Orthodontics, School of Dentistry, Faculty of Health Sciences, University of Pretoria, South Africa

²Dept. of Community Dentistry, School of Dentistry, Faculty of Health Sciences, University of Pretoria, South Africa



ARTICLE INFO

Article history:

Received 02-10-2023

Accepted 26-10-2023

Available online 16-01-2024

Keywords:

Malocclusion

Index of orthodontic treatment need

Dental health component

Waiting list

ABSTRACT

Background: The study described the patterns of malocclusion traits and the orthodontic treatment needs at a Dental Hospital using the Index of Orthodontic Treatment Needs (IOTN), Dental Health Component (DHC).

Materials and Methods: We evaluated 2079 pre-treatment study models and clinical records of orthodontic patients from the Department of Orthodontics waiting list. The assessment of malocclusion was measured according to the molar relationship using Angle's classification.

The DHC was further used to assess the five malocclusion traits: - missing teeth, overjet, crossbite, displacement of contact point, and overbite, including open bite (MOCCDO). The MOCCDO score was calculated to determine the orthodontic treatment needed based on the DHC grades 1–5. The data was analysed using SPSS version 28 and the level of significance was set at $p \leq 0.05$.

Results: The sample consisted of 59.3% (n=1232) females and 40.7% (n= 847) males. The mean age of the study sample was 14 years (SD±2.3) and ranged between 12 to 20 years. Angle's Class I malocclusion was found in 57.7% of the sample, followed by Class II (35.4%) and Class III (6.9%).

The orthodontic treatment needs DHC grade were: 53% grade 4 (need for orthodontic treatment), 21% grade 5 (great need for orthodontic treatment), 17% grade 3 (borderline need for orthodontic treatment), 6% grade 2 (little need for orthodontic treatment), and 3% grade 1 (no orthodontic treatment need).

A statistically significant difference was found in Angle Class I malocclusion and the DHC grades ($p=0.001$), as well as MOCCDO variables and DHC grades ($p=0.001$).

Conclusion: Angle's Class I malocclusion was the most predominant in our study sample compared to Class II and Class III malocclusions. The DHC of IOTN was successfully used to evaluate the severity of malocclusion with the majority of the sample in Grades 4 and 5, requiring mandatory orthodontic treatment. A high frequency of the occlusal traits included missing teeth, overjet, crossbite, displaced contacts and overbite. The results of our study showed that most patients on the orthodontic waiting list require mandatory orthodontic treatment. Therefore, it is recommended that DHC be used to place patients on the waiting list to prioritise orthodontic treatment.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Malocclusion is a developmental condition with a deflection from the normal relation or alignment of the teeth to other teeth in the same and opposing arch.¹ Malocclusion is

* Corresponding author.

E-mail address: phumzile.hlongwa@up.ac.za (P. Hlongwa).

ranked third among worldwide public health dental disease priorities after dental caries and periodontal disease.²

The prevalence of malocclusion varies from one country to another, between different gender and race, and with age.^{3–5} The worldwide prevalence of malocclusion among children and adolescents is 56% without relevant sex differences.⁶ The epidemiology of malocclusion among the continents has shown that Africa had the highest prevalence (81%), followed by Europe (71%), America (53%), and Asia (48%).⁶ The 2022 systemic review study reported a prevalence of 76.1% in Africa, however, South Africa was not included among the African countries.⁷

Epidemiology of malocclusion and assessment of orthodontic treatment needs are of national importance in many countries, and South Africa has included malocclusion epidemiology in the oral health national survey.⁸ The prevalence of malocclusion in South Africa has been reported to range from 32% to 52.3%.^{8–11}

An orthodontic treatment need index is an occlusal index originally developed to prioritize the need for orthodontic treatment. It classifies malocclusion and identifies patients in accordance with their treatment.^{12,13} Several indices for assessing malocclusions have been developed in the past,^{14–20} and the Index of Orthodontic Treatment Need (IOTN) has gained popularity.¹⁹ The IOTN is an epidemiological index used to estimate the need for orthodontic treatment and a screening instrument for determining treatment priority in government institutions with limited financial resources.²¹ The validity of the index has been assessed and confirmed by some researchers.^{22,23} It has also shown to be applicable in diverse ethnic groups without the need for modification,^{24,25} and was deemed to be trustworthy and easy to use.²¹

The IOTN was developed to grade malocclusion based on the significance of various occlusal traits for dental health and aesthetic impairment.¹⁹ The Dental Health Component (DHC) evaluates malocclusion utilising five occlusal traits: missing teeth, overjet, crossbite, displacement of the contact point, and overbite/open bite (MOCDO).²⁶ There are five grades of the DHC indicating orthodontic treatment need, grade 1 representing little or no need for treatment and grade 5 representing great need for treatment.¹⁹ The DHC malocclusion traits are further explained using alphabets to describe the severity of each malocclusion in the DHC grades (Table 1).²⁷

In South Africa's nine provinces, orthodontic treatment is performed mainly in private practice and the four dental teaching hospitals.²⁸ It has been reported that 80% of the population in South Africa depends on the government for health services, while only 20% can afford private care.²⁹ Thus the four dental teaching hospitals are inundated with orthodontic waiting lists for individuals with malocclusion. The treatment of malocclusion can be classified as interceptive orthodontic treatment; by contrast,

extensive care is performed in cases of moderate to severe malocclusion. Given this background of malocclusion in South Africa, it is necessary to evaluate the orthodontic treatment needs in relation to the severity of malocclusion. However, the prevalence of malocclusion and orthodontic treatment needs for patients on the orthodontic waiting list at a University Dental Hospital had not yet been reported in the indexed literature over the past 50 years. Therefore, this study assessed the orthodontic treatment needs based on the severity of malocclusion using the DHC of the IOTN. The results of the study may be used in future to plan sufficient treatment facilities and the development of adequate training programs for the prioritisation of orthodontic services and orthodontic waiting list reduction.

2. Materials and Methods

A cross-sectional retrospective study was undertaken utilising records of patients on the orthodontic waiting list at a University Dental Hospital. The study period was from 1 January 2012 to 31 December 2021. The dental hospital Ethical approval was obtained from the Research Ethics Committee (Reference: 22/2022) to conduct the study.

The study setting consisted of a University Dental Hospital situated in the city of Pretoria, in Gauteng Province, one of South Africa's nine provinces (Figure 1). Pretoria is the capital city of South Africa and was selected because the dental hospital provides oral health care, including orthodontic treatment, to most individuals in the South Africa's public health sector. Furthermore, the results of the study will generate new knowledge that will contribute to improvements in health care in the public health sector of South Africa.

In Gauteng Province, three university dental hospitals are available, separated by a radius of approximately 50 km. The fourth university dental hospital is situated in the Western Cape Province, about 1500 km away from Pretoria. All other seven provinces of South Africa do not have dental schools, and they rely on these four for public tertiary dental care including orthodontic treatment. Therefore, the University of Pretoria Dental Hospital location in the capital city of South Africa is easily accessible to the public and also serve as the referral centre for surrounding provinces and neighbouring countries.

The following inclusion and exclusion criteria were applied to the models to obtain the sample of the study:

2.1. Inclusion criteria

1. Patients in permanent dentition,
2. Age of 12 to 20 -years old,
3. Ideal orthodontics study models (correctly trimmed models with accurate reproducibility of the soft and hard tissue).



Figure 1: Map of South Africa with provinces and neighbouring countries 1- Limpopo Province, 2- Mpumalanga Province, 3- Gauteng Province, 4- North West Province, 5- Free State Province, 6- KwaZulu Natal Province, 7- Northern Cape Province, 8- Eastern Cape Province, 9- Western Cape Province
Source - https://en.wikipedia.org/wiki/Provinces_of_South_Africa

2.2. Exclusion criteria

1. Clefts and other craniofacial deformities.
2. Fractured models, restored teeth, and attrition.
3. Mixed dentition and primary dentition.
4. Models with no wax bite.
5. Significant periodontal disease.
6. Missing data, including age, biological sex and incomplete information.

2.3. Data collection

The assessment of dental classification of malocclusion was evaluated according to the molar relationship using Angle’s classification.³⁰ Furthermore, the severity of malocclusion was scored according to the DHC of the IOTN. The malocclusion measurement included the acronym MOCDO (Missing teeth; Overjet; Crossbites; Displacement of contact points; Overbite) to score the worst deviant occlusal trait of the malocclusion (Table 2). Missing teeth were counted, while other measurements were done using a transparent plastic ruler with a tip of 0 mm. The most severe measurement was used as the DHC score, which classified orthodontic treatment needs.

The principal investigator (TN) and one of the authors (MALM) were calibrated prior to the commencement of the study until the level of agreement (Kappa value = 0.9) was attained.

Repeated measurements of the 10% of the study sample were re-examined two weeks after the initial measurements to test for intra-examiner and inter-examiner reliability. The level of agreement attained for intra-examiner reliability was Kappa score = 0.8, and inter-examiner reliability Kappa score = 0.9.

The data was analysed using SPSS version 28, Chi-square test was used to test for association between Angle’s classification, MOCDO and DHC grades and the level of significance was set at $p \leq 0.05$.

3. Results

We measured 2079 records, which consisted of 59% (1232) females and 41% (847) males. The mean age of the study sample was 14 years (SD ± 2.3) and ranged between 12 to 20 years. Class I malocclusion was found in 58% of the sample, followed by Class II (35%) and Class III (7%). The majority of the sample (53%) had a great need for orthodontic treatment, indicated by the DHC grade 4, followed by grade 5 (21%) having the greatest need and grade 1 had the lowest proportion (3.2%) with no need for treatment as shown in Table 3.

Figure 2 illustrates the distribution of different malocclusion traits within the DHC grades according to the MOCDO acronym. The missing/unerupted permanent teeth within the study sample were observed in DHC 4h (n=121;5.8%) followed by DHC 5h (n=66;3.1%). The most predominant occlusal trait was the displacement of contacts found in DHC 4c (n=561;27%), followed by DHC 4t (451;21.7%). Overjet was mostly found in DHC 4a (n=275;13.2%), followed by DHC 5a (n=231;11.1%), with the least being DHC 2b (n=99;4.8%). Most anterior and posterior crossbites were in the DHC 3c (n=451; 21,7%), followed by DHC 4c (407;19%). Overbite was predominantly found in DHC 2f (297;14.3%) and DHC 3f (198;5%).




Figure 2: Distribution of malocclusion traits within the DHC scores

Table 4 presents the association between Angle’s classification, MOCDO and DHC grades. More patients

with Angle Class I malocclusion presented with a great need of treatment (grade 4) compared to Angle Class II and Class III, and the Chi-square test showed statistically significant differences ($p=0.001$). The missing teeth component of MOCDO also showed a Chi-square statistically significant difference ($p=0.001$), with most of the sample in grade 4 compared to grade 5. Overjet was found more in grade 4 compared to grade 5 and 3 with a Chi-square statistically significant difference ($p=0.001$). Most of the sample with crossbites, displaced contacts and overbites were in grade 4 compared to grade 5, 3 and 2, and the Chi-square showed a statistically significant difference ($p=0.001$).

Table 1: Alphabet representation of malocclusion in the DHC grades

Alphabets	Malocclusion description
a	Overjet – recorded to the most prominent part of the most prominent incisor
b	Reverse overjet with no masticatory or speech problems
c	Crossbite
d	Displacement of contact points where teeth deviate from the line of the arch, worst displacement recorded, spacing inline of the arch not included
e	Open bite
f	Deep bite
g	Good occlusion
h	Hypodontia
i	Impacted due to lack of space ≤ 4 mm
l	Posterior lingual crossbite
m	Reverse overjet with masticatory or speech problems
p	Defects of the cleft lip and palate
s	Submerged deciduous teeth
t	Partially erupted teeth, tipped and impacted against adjacent teeth
x	Presence of supernumerary teeth

4. Discussion

Our study has evaluated the prevalence of malocclusion from patients on a university dental hospital orthodontic waiting list and found a very high number with the greatest need for orthodontic treatment. The results were expected because the dental hospital is a referral centre for specialized orthodontic care from general dental practitioners and neighbouring clinics, provinces, and countries. Our study results showed that most of the sample were female (59%), similar to the results reported by Nugroho et al., who found that most of the recipients for orthodontic treatment were women.³¹ A study by Buthelezi et al. also revealed that most of the patients consulting at their institution were female.³² This could be explained by the fact that most females are conscious of their appearance.³³

The findings of our study showed that Angle Class I malocclusion (58%) was predominant compared to Class II and III. Similar results have been reported previously, showing Class I malocclusion predominancy followed by Class II and III, respectively.³⁴ The difference in the frequency of the various Angle's classes in most studies could be due to different geographic locations and ethnicity of the study sample. A 2020 systematic review and meta-analysis report on worldwide prevalence of malocclusion was only on four studies conducted in the continent of Africa (Tanzania 2009, Nigeria 2010, Morocco 2012 and Libya 2013). The results of the review were similar to our findings with that Angle Class I was predominant (61%), followed by Class II and Class III.⁶ Similarly, Yemitan and Oyapero reported that Class I malocclusion was the highest (76.7%), in their systematic review study in Africa, followed by Class II and Class III was the lowest among all Angle's classes of malocclusion.⁷ However, these systematic review studies were conducted in the Northern part of the continent of Africa and the countries reported are not a representation of the 54 countries in the continent, world's second-largest and second-most populous continent. Consequently, comparative studies in other African continent are still lacking especially, no studies from the Southern part of the continent were reported in these reviews.

Our study findings showed that most of the patients requiring orthodontic treatment needs were in DHC grade 4 (53%) with great need for treatment and grade 5 (21%) with the greatest need for orthodontic treatment. Similarly, to our findings, Maumela and Hlongwa reported that 21.7% of their sample had a severe need for orthodontic treatment, while 41.7% required mandatory orthodontic treatment.³⁵ The distribution of malocclusion severity in our study was similar to the findings of Shue-Te Yeh et al., who reported that the majority of their patient presented with DHC grade 4.³⁶ Our results found 17% of patients with a moderate need for treatment (grade 3) and 6% with little need for treatment (grade 2) while 3%, had no need for treatment (grade 1). The inability of these grades to allocate the patients for orthodontic treatment does not mean that they do not present with malocclusion, but their malocclusion fell below the category of treatment deemed by the DHC treatment needs.

The missing/unerupted component of the MOCDO in our study was 9% ($n=187$). Similar findings have been reported on missing teeth ranging from 2.6% to 11.3%.³⁷ Contrary to our findings, Mani et al. reported a lower percentage of 3.2% of missing teeth in their study.³⁸ Our study did not explore the reasons for the missing teeth in our sample, but these could include; early tooth loss due to extraction, dental caries or congenitally missing.

The presence of overjet in our study sample was 55%. This is a very important factor to consider, as patients with an increased overjet are said to have an increased risk of trauma to the anterior teeth.³⁹ Lower prevalence

Table 2: DHC components, using the MOCDO acronym

IOTN Dental Health Component	Grade 5	Grade 4	Grade 3	Grade 2	Grade 1
Missing teeth	5h = extensive hypodontia + restorative implications >1 tooth missing per quadrant requiring pre-restorative orthodontic treatment 5s = submerging primary teeth 5i = impeded eruption/impaction	4h = less extensive hypodontia requiring orthodontic treatment for pre-restorative or space closure			
Overjet		4a = OJ 6.1-9 mm 4b = ROJ >3.5 mm with no masticatory and speech difficulties	3a = OJ 3.6-6 mm + incompetent lips 3b = ROJ 1.1-3.5 mm	2a = OJ 3.6-6 mm + competent lips 2b = ROJ 0.1-1 mm	
Crossbite		4c = x-bites + >2 mm discrepancy between RCP and ICP 4l = posterior lingual x-bite	3c = x-bite + 1.1-2 mm discrepancy between RCP and ICP	2c = x-bite with up to 1 mm discrepancy between ICP and RCP	
Displacement of contact point		4d = contact point displacement >4 mm 4t = partially erupted teeth, tipped and impacted against adjacent teeth 4x = supplemental teeth	3d = contact point displacement 2.1-4 mm	2d = contact point displacement 1.1-2 mm	Minor irregularity
Overbite		4e = lateral or anterior open bite >4 mm 4f = increased + complete OB + gingival or palatal trauma	3e = lateral or anterior open bite 2.1-4 mm 3f = increased + complete OB with no gingival trauma	2e = lateral or anterior open bite 1.1-2 mm 2f = increased OB >3.5 mm and no gingival contact	

Source²⁷

Table 3: Demographics characteristics of the study sample

Variables	Frequency
Mean age in years (SD)	14(±2.34)
Age range in years	12 - 20
Gender	n (%)
Male	847 (41%)
Female	1232 (59%)
Angle's Classification	
Class 1	1199 (58%)
Class 2	737 (35%)
Class 3	143 (7%)
DHC	
Grade 1	66 (3%)
Grade 2	132 (6%)
Grade 3	341 (17%)
Grade 4	1100 (53%)
Grade 5	440 (21%)

Table 4: Association between Angle's Classification, DHC malocclusion traits and DHC categories

	DHC Grades					p-value
	1	2	3	4	5	
Angle Class I	55(2.6%)	99(4.8%)	264(12.7%)	583(28%)	198(9.5%)	0.001
Angle Class II	11(0.5%)	22(1%)	55(2.6%)	462(22.2)	187(8.9)	
Angle Class III	0(0%)	11(0.5%)	22(1%)	55(2.6%)	55(2.6%)	0.001
Missing Teeth	0(0%)	0(0%)	0(0%)	121(5.8%)	66(3.2%)	
Overjet	0(0%)	0(0%)	154(7.3%)	649(31%)	330(15.9%)	0.001
Crossbite	0(0%)	55(2.6%)	143(6.9%)	704(33.9)	264(12.7%)	0.001
Displaced contacts	0(0%)	99(4.8%)	319(15.3%)	1001(48.1%)	396(19%)	0.000
Overbite	0(0%)	22(1.1%)	132(6.3%)	484(23.3%)	220(10.5%)	0.001

of overjet compared to our study have been reported by Singh et al. with a finding of 17.5%.⁴⁰ A study on Nigerian subjects also showed a lower prevalence of 43% overjet, compared to our study findings.⁴¹ Research conducted on the female population of Saudi Arabia seeking orthodontic treatment found 76% overjet in adolescents which was higher compared to our results.⁴²

The anterior and posterior crossbite in our sample was found to be 56.1%, similarly to the study reported by Albarakari et al. who also found a similar prevalence of crossbites in 60.5% of their study.⁴² Higher prevalence of crossbite compared to our study was reported by Gungor et al. in 64.3% of their patients⁴³ However, Shrestha et al.⁴⁴ reported a lower prevalence of 23.3% in their study compared to our findings. In our study, anterior crossbites (49.2%) were more common than posterior crossbites (6.9%). Our results were similar to a study by Ajayi et al. findings of 21.4% for the anterior crossbite and 12.2% for the posterior crossbite.⁴³ The reported prevalence for the posterior crossbite is low because patients do not seek orthodontic treatment for the correction of posterior teeth since they do not see or recognize it as a problem. However, it is important to diagnose and manage posterior crossbites as they may be associated with symptoms of the temporomandibular joint dysfunction.⁴⁵

Measurement of contact point displacement as the DHC component of IOTN was performed on the greatest displacement of point of contact. Clinically, displacement of contact points is also called crowded teeth. In our study, displacement of points of contact of 87% (n= 1815) was the most predominant malocclusion trait. Similar results have been reported on crowding as the common type of malocclusion.^{40,46}

The measurement of overbite was found to range from 3.5mm to overbite covering the whole lower incisors, with or without palatal trauma. Our study had a prevalence of 29.6% of overbite comparable to the findings of 20.7% of Nepalese patients⁴¹ and 25% of Nigerian patients.^{41,44} Our study findings showed that 14.3% of the sample had an overbite greater than 3.5mm, with 9.5% having increased overbite without palatal contact, while 5.8% had an increased overbite with palatal trauma. Contrary to

our findings, Souames et al. found an increased overbite prevalence of 15%, an overbite without palatal contact of 45%, and an overbite with palatal contact of 10% in their samples.⁵

The results of our study found that the percentage of patients with open bites was 11.7%. Previous findings have demonstrated a reduced prevalence of 8%.^{41,44} In contrast, the prevalence of open bite was greater in a sample of Saudi patients (46.7%),⁴² compared to our study findings. The variation between prevalence of overbite (29.6%) and open bites (11.7%) in our study sample is similar to a previous study that reported that overbite (13.23%) was more predominant than open bites (2%).⁴⁰

5. Study Limitations

Our study experienced challenges with incomplete records, due to the nature of retrospective investigations. The study records that were evaluated are not collected for research, however, the challenge was addressed by adhering to the inclusion and exclusion criteria for the sample selection. The skeletal malocclusion patterns of the samples were also not evaluated for this study because lateral cephalometric radiographs were not assessed. Furthermore, our sample had more females compared to males, and this cannot be generalized for biologic sex differences. The study findings showed the patients that had no need for treatment categorized by grade 1 of the DHC index showed that the index was not sensitive to patients with mild malocclusion. This exclusion by the index does not imply that those patients do not have malocclusion requiring treatment, but the index did not categorise them for treatment for government institutions with limited funds.

6. Conclusion

Angle's Class I malocclusion was the most prevalent in our study sample compared to Class II and Class III malocclusions. The DHC was successfully used to evaluate the severity of malocclusion with the majority of the sample in Grades 4 and 5, requiring mandatory orthodontic treatment. A high frequency of the occlusal traits included missing teeth, overjet, crossbite, displaced contacts and

overbite.

7. Recommendations

The results of our study showed that most patients on the orthodontic waiting list require mandatory orthodontic treatment. Therefore, it is recommended that DHC be used for placing patients on the waiting list in government funded institutions to ensure that patients with severe malocclusion are prioritised for treatment. Multicentre studies on the prevalence of malocclusion should be firstly conducted in the country of South Africa and also across the continent of Africa for clinicians and policy makers to develop strategies to prevent and manage this anomaly.

8. Data Availability Statement

The data supporting this study's findings are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

9. Ethics Approval Statement

Ethical approval was obtained from the University Research Ethics Committee (Reference: 22/2022) to conduct the study.

10. Source of Funding

None.

11. Conflict of Interest

None.


References


- Baskaradoss J, Bhagavatula P. Measurement and distribution of malocclusion, trauma, and congenital anomalies. In: Burt and Eklund's Dentistry, Dental Practice, and the Community-E-Book. Philadelphia: WB Saunders Company; 2020. p. 208–17. doi:10.1046/j..2003.com122.x.
- Petersen PE. The World Oral Health Report 2003: continuous improvement of oral health in the 21st century—the approach of the WHO Global Oral Health Programme. *Community Dent Oral Epidemiol.* 2003;31(Suppl 1):3–23. Available from: <https://doi.org/10.1046/j..2003.com122.x>.
- Mtaya M, Brudvik P, Åström AN. Prevalence of malocclusion and its relationship with socio-demographic factors, dental caries, and oral hygiene in 12- to 14-year-old Tanzanian schoolchildren. *Eur J Orthod.* 2009;31(5):467–76.
- Al-Zubair NM. Orthodontic treatment need of yemeni children assessed with dental aesthetic index. *J Orthod Sci.* 2014;3(2):41–5.
- Souames M, Bassigny F, Zenati N, Riordan PJ, Boy-Lefevre ML. Orthodontic treatment need in french schoolchildren: An epidemiological study using the index of orthodontic treatment need. *Eur J Orthod.* 2006;28(6):605–9.
- Lombardo G, Vena F, Negri P, Pagano S, Barilotti C, Paglia L, et al. Worldwide prevalence of malocclusion in the different stages of dentition: A systematic review and meta-analysis. *Eur J Paediatr Dent.* 2020;21(2):115–22.
- Yemitan TA, Oyapero AO. Prevalence of malocclusion in africa: A systematic review and meta-analysis. *Magna Sci Adv Res Rev.* 2022;5(1):30–5.
- Wyk PJV, Wyk CV. Oral health in south africa. *Int Dent J.* 2004;54(6 Suppl 1):373–7.
- Wyk PV, Drummond RJ. Orthodontic status and treatment need of 12-year-old children in south africa using the dental aesthetic index. *S Afr Dent J.* 2005;60(8):334–6.
- Hlongwa P. Malocclusion among 12-year-old school children in mankweng, limpopo province of south africa. *S Afr Dent J.* 2005;60(10):455–7.
- Booyesen JL, Harris A. Orthodontic treatment need and demand in the upington area of the northern cape province. South Africa; 2017. Available from: https://etd.uwc.ac.za/bitstream/handle/11394/6061/booyesen_msc_dnt_2017.pdf?sequence=1&isAllowed=y.
- Tang EL, Wei SH. Recording and measuring malocclusion: A review of the literature. *Am J Orthod Dentofacial Orthop.* 1993;103(4):344–51.
- Carlos JP. Evaluation of indices of malocclusion. *Int Dent J.* 1970;20(4):606–17.
- Summers CJ. The occlusal index: A system for identifying and scoring occlusal disorders. *Am J Orthod.* 1971;59(6):552–67.
- Massler M, Frankel JM. Prevalence of malocclusion in children aged 14 to 18 years. *Am J Orthod.* 1951;37(10):751–68.
- Kirk LEV, Pennell EH. Assessment of malocclusion in population groups. *Am J Public Health Nations Health.* 1959;49(9):1157–63.
- Draker HL. Handicapping labio-lingual deviations: A proposed index for public health purposes. *Am J Orthod.* 1960;46(4):295–305.
- Salzmann J. Handicapping malocclusion assessment to establish treatment priority. *Am J Orthod.* 1968;54(10):749–65.
- Brook PH, Shaw WC. The development of an index of orthodontic treatment priority. *Eur J Orthod.* 1989;11(3):309–20.
- Grainger R. Orthodontic treatment priority index. *Vital Health Stat 2.* 1967;(25):1–49.
- Hlongwa P, Beane R, Seedat A, Owen C. Orthodontic treatment needs: Comparison of two indices. *South Afr Dent J.* 2004;59(10):421–4.
- Burden DJ, Pine CM, Burnside G. Modified iotn: An orthodontic treatment need index for use in oral health surveys. *Community Dent Oral Epidemiol.* 2001;29(3):220–5.
- Siddiqui TA, Shaikh A, Fida M. Agreement between orthodontist and patient perception using index of orthodontic treatment need. *Saudi Dent J.* 2014;26(4):156–65.
- Ahmed B, Gilthorpe MS, Bedi R. Agreement between normative and perceived orthodontic need amongst deprived multiethnic school children in london. *Clin Orthod Res.* 2001;4(2):65–71.
- Dawjee S, Ackerman A, Shaw W. An aesthetic component of the iotn for black subjects. *South Afr Dent J.* 2002;57(7):258–63.
- Richmond S, O'Brien K, Burden BI. An introduction to occlusal indices. UK: Mandent Press; 1994.
- Jawad Z, Bates C, Hodge T. Who needs orthodontic treatment? Who gets it? And who wants it? *British Dental Journal.* 2015;218(3):99–103.
- De Müelenaere J, Wiltshire W. Orthodontic treatment in general dental practice in South Africa. *J Dent Assoc S Afr.* 1990;45(12):529–33.
- Coovadia H, Jewkes R, Barron P, Sanders D, Mcintyre D. The health and health system of south africa: Historical roots of current public health challenges. *Lancet.* 2009;374(9692):817–34.
- Angle EH. Classification of malocclusion. *Dent Cosmos.* 1899;41:350–75.
- Nugroho MJ, Ismah N, Purbiati M. Orthodontic treatment need assessed by malocclusion severity using the dental health component of iotn. *J Int Dent Med Res.* 2019;12(3):1042–6.
- Buthlezi NL, Madiba TK. Oral hygiene habits and status of orthodontic patients attending the university of pretoria, oral and dental hospital. *South Afr Dent J.* 2021;76(3):130–5.
- Tin-Oo MM, Saddki N, Hassan N. Factors influencing patient satisfaction with dental appearance and treatments they desire to improve aesthetics. *BMC Oral Health.* 2011;11:1–8.
- Perrotta S, Bucci R, Simeon V, Martina S, Michelotti A. Prevalence of malocclusion, oral parafunctions and temporomandibular disorder-pain in italian schoolchildren: An epidemiological study. *J Oral*

- Rehabil.* 2019;46(7):611–6.
35. Maumela PM, Hlongwa P. Application of the dental aesthetic index in the prioritisation of orthodontic service needs: Clinical review. *SADJ.* 2012;67(7):380–3.
 36. Yeh MST, Koochek AR, Vlaskalic V, Boyd R, Richmond S. The relationship of 2 professional occlusal indexes with patients' perceptions of aesthetics, function, speech, and orthodontic treatment need. *Am J Orthod Dentofacial Orthop.* 2000;118(4):421–8.
 37. Larmour CJ, Mossey PA, Thind BS, Forgie AH, Stirrups DR. Hypodontia-a retrospective review of prevalence and etiology. Part i. *Quintessence Int.* 2005;36(4):263–70.
 38. Mani SA, Mohsin WSY, John J. Prevalence and patterns of tooth agenesis among malay children. *Southeast Asian J Trop Med Public Health.* 2014;45(2):490–8.
 39. O'Brien K, Wright J, Conboy F, Chadwick S, Connolly I, Cook P, et al. Effectiveness of early orthodontic treatment with the twin-block appliance: A multicenter, randomized, controlled trial. Part 2: Psychosocial effects. *Am J Orthod Dentofacial Orthop.* 2003;124(5):488–94.
 40. Singh VP, Sharma A. Epidemiology of Malocclusion and Assessment of Orthodontic Treatment Need for Nepalese Children. *Int Sch Res Notices.* 2014;2014:768357.
 41. Ajayi E, Ize-Iyamu I. Malocclusion and occlusal traits among orthodontic patients seen at the university of benin teaching hospital, nigeria. *Niger Dent J.* 2010;18(1):12–6.
 42. Albarakati SF, Taher S. Malocclusion traits in saudi females seeking orthodontic treatment. *Pak Oral Dent J.* 2010;30(1):127–32.
 43. Gungor K, Taner L, Kaygisiz E. Prevalence of posterior crossbite for orthodontic treatment timing. *J Clin Pediatr Dent.* 2016;40(5):422–4.
 44. Shrestha S, Shrestha RM. An analysis of malocclusion and occlusal characteristics in nepalese orthodontic patients. *Orthodontic Journal of Nepal.* 2013;3(1):19–25.
 45. Thilander B, Bjerklin K. Posterior crossbite and temporomandibular disorders (TMDs): need for orthodontic treatment? *Eur J Orthod.* 2012;34(6):667–73.
 46. Al-Emran S, Wisth PJ, Böe OE. Prevalence of malocclusion and need for orthodontic treatment in saudi arabia. *Community Dent Oral Epidemiol.* 1990;18(5):253–5.


Author biography

Tshepiso D Ntseke, -  <https://orcid.org/0000-0002-9675-0409>

Thomas K Madiba, -  <https://orcid.org/0000-0002-0171-0595>

Millicent Motsepe, -  <https://orcid.org/0000-0002-1346-6681>

Mpule AL Moshaoa, -  <https://orcid.org/0000-0003-0306-3739>

Phumzile Hlongwa, -  <https://orcid.org/0000-0002-8052-9275>

Cite this article: Ntseke TD, Madiba TK, Motsepe M, Moshaoa MAL, Hlongwa P. The distribution of malocclusion using the index of orthodontic treatment needs at a university dental hospital in and around Pretoria, South Africa. *Int J Oral Health Dent* 2023;9(4):258-265.