

An investigation into the potential use of Dennkur™ as an intra-canal irrigant and medicament in Endodontics

Sumita Bhagwat^{1,*}, Amrut Bambawale², Anchal Mehta³, Leena Padhye⁴

¹Professor, ²⁻³Senior Registrar, ⁴Professor & HOD, Dept. of Conservative Dentistry, Dr. DY Patil University School of Dentistry, Nerul, Navi Mumbai

***Corresponding Author:**

Email: sumital572@gmail.com

Abstract

Achieving complete asepsis within the root canals during Endodontic therapy is an elusive goal of every dental practitioner. While many commercial root canal irrigants and medicaments are employed to achieve the same, they are not without undesirable side effects. A fair amount of scientific interest is currently being generated by the potential of traditional herbal agents and their use in achieving asepsis in the root canal. Research in nearly all branches of medicine exists documenting the potential of Turmeric as an antimicrobial agent. Curcumin, the active ingredient in Turmeric, has shown fair results in its antimicrobial action in Periodontics. Our in vitro study explores the antimicrobial potential of a curcumin-containing product Dennkur™ against endodontic pathogens *E.faecalis* and *C.albicans* by the Agar cup method.

Keywords: Curcumin, Dennkur, *E.faecalis*, *C.albicans*

Introduction

The success of endodontic treatment depends on the eradication of microorganisms from the root-canal system and prevention of reinfection. The root canal is shaped with hand and rotary instruments under constant irrigation to remove inflamed and necrotic tissue, microbes/biofilms, and other debris from the root-canal space. Several studies using advanced techniques such as microcomputed tomography (CT) scanning have demonstrated that proportionally large areas of the main root-canal wall remain untouched by the instruments,⁽¹⁾ emphasizing the importance of chemical means of cleaning and disinfecting all areas of the root canal. Presently the most reliably and universally used root canal irrigants in endodontic practice are Sodium hypochlorite and Chlorhexidine gluconate, Calcium hydroxide is considered as a gold standard for use as an intracanal medicament in endodontic cases with suspected infection.

Due to the clinical drawbacks encountered with the use of the above irrigants and medicaments, research has shifted into the direction of traditional medicine in search for an ideal irrigant and medicament. The rhizome Turmeric (*Curcuma longa*) is one such traditional asian herb that has sparked a great deal of interest among researchers.⁽²⁾ The active ingredient in Turmeric is Curcumin. Curcumin's potent anti-inflammatory properties have lead to active research on its use for a variety of inflammatory conditions, including postoperative inflammation, arthritis, uveitis, inflammatory pseudotumors, dyspepsia, irritable bowel syndrome, inflammatory bowel disease, pancreatitis, and *Helicobacter pylori* infection. Most studies are promising and further exploration of curcumin's therapeutic value for inflammatory conditions is warranted (Jurenka, 2009).^(3,4)

Dennkur™ is a mouth-dissolving pastille that contains Curcumin with its natural antiseptic, antibacterial and anti-inflammatory properties. It slowly dissolves in mouth giving a longer contact time for the local action on the floor of mouth. The natural antibacterial action of curcumin reduces the bacterial load in the oral cavity. Each soft gelatine pastille (Lozenge) contains *Curcuma Longa* (*Curcuma*) -100mg, excipients and flavouring agents.

Dennkur™ is believed to have the same antibacterial action as chlorhexidine and hence an investigation into its potential use as an endodontic irrigant and intracanal medicament was carried out. The aim of this study was to test the antibacterial action of Dennkur™ against endodontic pathogens *Enterococcus faecalis* and *Candida albicans*.

Materials and Method

Group 1 was Sterile Nutrient Agar (for *Enterococcus Faecalis*).

Group 2 was Sterile Sabourauds Agar (For *Candida Albicans*)

The following five samples were compared in the following in vitro study

1. 3% Sodium Hypochlorite (Prime Dental Products)
2. 2% Chlorhexidine (Stedman Pharmaceuticals Pvt Ltd)
3. Calcium Hydroxide (Prime Dental Products)
4. Curcumin (Dennkur™ Lozenges (1 tablet in 10 mL distilled water)
5. Normal saline

Media used were:

1. Sterile Nutrient Agar (for *Enterococcus Faecalis*)
2. Sterile Sabourauds Agar (For *Candida Albicans*)

Microbial cultures used were:

1. *Enterococcus faecalis* (NCIM no 5025)

2. *Candida albicans* (ATCC No. 10231)

Armamentarium used:

1. Petri plate
2. Agar
3. Cork boner

Procedure: The method employed to carry out the following in-vitro study was Agar Cup Method. The equipment was sterilized and a sterile environment in order to prevent cross contamination. Preparation of the culture mediums, i.e. Sterile Nutrient Agar (for *Enterococcus Faecalis*) and Sterile Sabourauds Agar (for *Candida Albicans*) was carried out and it was kept for boiling to maintain asepsis. After appropriate temperature was reached they were allowed to cool down to luke warm. The culture medium (Sterile Nutrient Agar) was removed by the use of pipette and was placed between two Bunsen burners, this was done to maintain a zone of absolute sterilization. 25 ml of sterile culture medium with an addition of 1 mL of cultural strain (Group I containing *Enterococcus Faecalis*) was poured in a petri plate and allowed to solidify to room temperature.

The same procedure was followed for second culture medium (Sterile Sabouraud's Agar) which was mixed with Group II strain (containing *Candida Albicans*) and poured into sterile petriplates and was allowed to cool at room temperature. The cork boner was placed in the aseptic zone between two Bunsen burners and 5 wells of 8 mm diameter were punched. After the wells were made, 0.1 mL of undiluted sample was introduced in the respective well.

Saline was used as a control. After the incubation period of 24hrs, the plates were removed and zones of inhibition were recorded. The experiment was repeated 3 times (each time after a gap of 48hrs) and zones of inhibition were recorded using Vernier Calipers in mm (millimeters).




Fig. 1: Action of Dennkur, calcium Hydroxide, chlorhexidine sodium hypochlorite on *E. faecalis* (Group 1)




Fig. 2: Zones of inhibition demonstrated on action with *E. faecalis* (Group 1)




Fig. 3: Action of Dennkur, calcium hydroxide, sodium hypochlorite, chlorhexidine on *Candida Albicans* (Group 2)




Fig. 4: Zones of inhibition demonstrated on action with *C. Albicans* (Group 2)

Results

Table 1: Enterococcus Faecalis on Nutrient Agar Plate (Group 1) - Zone of Inhibition (mm)

Materials	Undiluted	1:2 dilution	1:10 dilution
Sodium, Hypochlorite 3%	13mm	11.5mm	8mm
Calcium Hydroxide	13mm	12mm	0
Chlorhexidine 2%	21mm	18mm	18mm
Dennkur™	0	0	0
Saline	0	0	0

Table 2: Candida Albicans on a Sabouraud's Agar plate (Group 2) Zone of Inhibition (mm)

Materials	Undiluted	1:2 dilution	1:10 dilution
Sodium, Hypochlorite 3%	16mm	13mm	0
Calcium Hydroxide	14mm	12mm	0
Chlorhexidine 2%	15mm	12mm	12mm
Dennkur™	0	0	0
Saline	0	0	0

Discussion

By virtue of their mechanism of action, present day root canal irrigants and medicaments nearly effectively clean the root canal. They are able to disinfect and penetrate dentin and its tubules, offer long-term antibacterial effect (substantivity), remove the smear layer, and should be non-antigenic, nontoxic and non-carcinogenic. In addition, they require to have no adverse effects on dentin or the sealing ability of filling materials. Furthermore, they should be relatively inexpensive, convenient to apply and cause no tooth discoloration. The irrigants currently used that are considered gold standard are Sodium Hypochlorite & Chlorhexidine. The universally used and accepted intra-canal medicament is Calcium Hydroxide.

The most effective method for eliminating *E.faecalis* from the root canal space and dentinal tubules is by the use of Sodium hypochlorite (NaOCl) and Chlorhexidine in a gel or liquid concentration form. But the disadvantages of sodium hypochlorite are its unpleasant taste, toxicity, and its use is suspected in weakening of the tooth structure by decreasing the hardness and structural integrity of the dentin within the root canal.⁽⁵⁾

Chlorhexidine gluconate is a cationic bisguanide that seems to act by adsorbing onto the cell wall of the microorganism and causing leakage of intercellular

components. At low concentration, it has a bacteriostatic effect and at higher concentrations it has a bactericidal effects. It has a lower cytotoxicity as compared to sodium hypochlorite. However, unlike NaOCl, it cannot dissolve organic substances and necrotic tissues present in the root canal system. In addition, it is unable to kill all bacteria and is not effective in removing the smear layer as well. No adverse effects have been published regarding Chlorhexidine being used as an irrigant or intra-canal medicament. However, the direct effect in an *in vitro* test on human stem cells of apical papilla showed lack of viable cells after its use to remove the smear layer. The disadvantages of hypochlorite and calcium hydroxide have motivated researchers to explore the potential of different chemicals and herbs as root canal irrigants.⁽⁶⁾

Calcium hydroxide was originally introduced to the field of endodontics by Herman in 1920 as a pulp-capping agent, it was found effective as intra-canal medicament.⁽⁷⁾ The antimicrobial activity of Ca(OH)₂ is related to the release of hydroxyl ions in an aqueous environment (Siqueira 2001). Hydroxyl ions are highly oxidant free radicals that show extreme reactivity with several biomolecule. The lethal effects of hydroxyl ions on bacterial cells are probably due to (Siqueira & Lopes 1999) damage to the bacterial cytoplasmic membrane, protein denaturation and damage to the DNA.

The major disadvantage of Calcium hydroxide is its ineffectiveness in completely eliminating microorganisms. Various studies done by DiFiore et al. (1983), Haapasalo & Orstavik (1987), Safavi et al. (1990) have shown the ineffectiveness of Calcium hydroxide against *E.Faecalis* and *C.Albicans*.

Thus, looking at the various disadvantages and adverse effects of the mentioned irrigants and medicaments, there is a need for alternatives that would endeavor to eliminate residual microorganisms without causing any adverse effects. Various Indian herbal and ayurvedic products such as extracts from Neem (*Azadiracta Indica*), Ashwaghandha (*Withania Somnifera*) and Curcumin (*Curcuma Longa*) are being investigated as alternatives.

Turmeric or Curcumin with a chemical formula (1,7-bis(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione) and other curcuminoids constitute the main phytochemicals of *curcuma longa* L. (*Zingiberaceae* family) rhizome with common name of turmeric. Several Studies have reported the broad spectrum antimicrobial activity for curcumin including antibacterial, antiviral, antifungal and antimalarial activities.^(2,3,4,6,7,8) An *in vitro* study evaluated the efficacy of three intracanal medicaments 10% and 20% of turmeric, calcium hydroxide, 1% chlorhexidine gel and 1% chlorhexidine gel with 10% calcium hydroxide against *E.faeclis* in necrotic primary teeth and concluded that chlorhexidine with or without calcium

hydroxide was more effective than calcium hydroxide alone (Kumar, Hemanshi. 2013).

Neelakantan *et al* found that curcumin had significant antibacterial activity against *E.faecalis*.⁽⁹⁾ Haukyik *et al* studied the phototoxic effects of curcumin against gram positive bacteria like *E. faecalis*, streptococcus intermedius and gram negative bacteria *E. coli* were investigated in aqueous preparations.⁽¹⁰⁾

Prasanna *et al.*, (2011) conducted an *in vitro* study to evaluate the antimicrobial efficacy of curcumin against *E. faecalis*, *C.Albicans* considering sodium hypochlorite (3%) as reference for comparison. The result of his study revealed that curcumin had significant antibacterial activity against *E. faecalis* he concluded that the antibacterial activity of curcumin was similar to sodium hypochlorite and thus herbal medicine can be used in endodontics for root canal failure.⁽¹¹⁾

Lívia N. Dovigo, Ana Cláudia Pavarina, Ana Paula D. Ribeiro (2011) also carried out an *in vitro* investigation and found that Curcumin was effective against *Candida Albicans*.⁽¹²⁾

But Hegde and Kesaria⁽¹³⁾ found that Turmeric showed no antibacterial action against *E.Faecalis* in an *in vitro* study by agar cup method testing antibacterial action of several new agents for use as root canal irrigants.

In our study, we observed that the antibacterial action of the test sample was nil in all dilutions, the readings for the test sample, sodium hypochlorite, chlorhexidine, calcium hydroxide were tabulated and statistically analysed by ANOVA showed the results highly significance ($p > 0.05$).

The active component of Dennkur™ (Curcumin) was claimed to have antiseptic as well an anti-inflammatory activity. It was also suggested to have endodontic use and claimed to be as effective as chlorhexidine. Since no specific study was done to show the effectiveness of the product (Dennkur™) against endodontic flora, the present study was carried out to investigate the antibacterial activity of this product against *E.Faecalis* and *C.Albicans*. In our study, Dennkur™ showed zero zone of inhibition against *E faecalis* & *C.Albicans* as compared with the gold standard irrigants and medicament, which showed zones of inhibition ranging from 8-16mm (as shown in Table 1 and Table 2).

Conclusion

Curcumin has shown a good potential in terms of antibacterial activity to warrant further research. However the commercial product Dennkur™ in its present formulation appears to be ineffective in showing the similar antibacterial activity. While Dennkur™ shows use as oral lozenges, its activity against endodontic pathogens seems to indicate that this commercial product in its present formulation does not have any endodontic application.

Disclaimer

The authors state that there is no conflict of interest and no funding was received from any body to carry out this original research.

References

1. Markus Haapasalo, Ya Shen, Wei Qian, Yuan Gao. Irrigation in Endodontics Dent Clin N Am 54 (2010) 291-312.
2. Rana Pratap Singh, D. A. Jain. Evaluation of antimicrobial activity of curcuminoids isolated from turmeric. Int. J. of Pharm. & Life Sci., Vol. 3, Issue 1:Jan.:2012, 1368-1376.
3. Soheil Zorofchian Moghadamtousi, Habsah Abdul Kadir, Pouya Hassandarvish, Hassan Tajik, Szalay Abubakar and Keivan Zandi. A Review on Antibacterial, Antiviral, and Antifungal Activity of Curcumin. BioMed Research International Volume 2014 (2014), Article ID 186864, 12 pages.
4. Praveenkumar S Mandroli, Kishor Bhat. An *in vitro* evaluation of antibacterial activity of curcumin against common endodontic bacteria. Journal of Applied Pharmaceutical Science Vol. 3 (10), pp. 106-108, October, 2013.
5. Ravishankar. P, Lakshmi. T, Aravind K.S. Ethno-Botanical Approach for Root Canal Treatment -An Update. J. Pharm. Sci. & Res. 2011 Vol.3 (10), 1511-151.
6. Reshma Suvama, Sham S Bhat, K Sandeep Hedge. Antibacterial activity of Turmeric against *E.faecalis* - An *In vitro* study. International Journal Of Current Microbiology And Applied Sciences 2014, Vol3(2): 498-504.
7. Mohammadi Z, Soltani MK, Shalavi S. An update on the management of endodontic biofilms using root canal irrigants and medicaments. Iran Endod J. 2014 Spring;9(2):89-97. Epub 2014 Mar 8.
8. Chaturvedi TP. Uses of turmeric in dentistry: An update. Indian J Dent Res 2009;20:107-9.
9. Neelakantan P, Subbarao C, Sharma S, Subbarao CV, Garcia-Godoy F, Gutmann JL. Effectiveness of Curcumin against *Enterococcus faecalis* biofilm. Acta Odontol Scand. 2013 Nov;71(6):1453-7.
10. Haukvik T1, Bruzell E, Kristensen S, Tønnesen HH. Photokilling of bacteria by curcumin in selected polyethylene glycol 400 (PEG 400) preparations. Studies on curcumin and curcuminoids, XLI. Pharmazie. 2010 Aug;65(8):600-6.
11. Prasanna Neelakantan, Nithya Jagannathan, Nabeel Nazar. Ethnopharmacological approach in Endodontic Treatment: A Focused review. Int. J. of Drug Dev. & Res., Oct-Dec 2011, 3(4): 68-77.
12. Dovigo LN, Carmello JC, de Souza Costa CA, Vergani CE, Brunetti IL, Bagnato VS, Pavarina AC. Curcumin-mediated photodynamic inactivation of *Candida albicans* in a murine model of oral candidiasis. Med Mycol. 2013 Apr;51(3):243-51.
13. Hegde V., Kesaria D. Comparative evaluation of antimicrobial activity of neem, propolis, turmeric, liquorice and sodium hypochlorite as root canal irrigants against *e. Faecalis* and *c. Albicans* - an *in vitro* study. Endodontology 2013 Dec;25(2):38-45.